arXiv:1601.01779v1 [math.AC] 8 Jan 2016

ON FUNCTION COMPOSITIONS THAT ARE POLYNOMIALS

ERHARD AICHINGER

ABSTRACT. For a polynomial map $\boldsymbol{f} : k^n \to k^m$ (k a field), we investigate those polynomials $g \in k[t_1, \dots, t_n]$ that can be written as a composition $g = h \circ \boldsymbol{f}$, where $h : k^m \to k$ is an arbitrary function. In the case that k is algebraically closed of characteristic 0 and $\boldsymbol{f}$ is surjective, we will show that $g = h \circ \boldsymbol{f}$ implies that h is a polynomial.

1. INTRODUCTION

In the present note we investigate the situation that the value of a polynomial depends only on the value of certain given polynomials. To be precise, let k be a field, $m, n \in \mathbb{N}$, and let $g, f_1, \dots, f_m \in k[t_1, \dots, t_n]$. We say that g *is determined by* $\boldsymbol{f} = (f_1, \dots, f_m)$ if for all $\boldsymbol{a}, \boldsymbol{b} \in k^n$ with $f_1(\boldsymbol{a}) = f_1(\boldsymbol{b})$, $\dots$, $f_m(\boldsymbol{a}) = f_m(\boldsymbol{b})$, we have $g(\boldsymbol{a}) = g(\boldsymbol{b})$. In other words, g is determined by $\boldsymbol{f}$ if and only if there is a function $h : k^m \to k$ such that

$$g(\boldsymbol{a}) = h(f_1(\boldsymbol{a}), \dots, f_m(\boldsymbol{a})) \text{ for all } \boldsymbol{a} \in k^n.$$

For given $f_1, \dots, f_m \in k[t_1, \dots, t_n]$, the set of all elements of $k[t_1, \dots, t_n]$ that are determined by $(f_1, \dots, f_m)$ is a k-subalgebra of $k[t_1, \dots, t_n]$; we will denote this k-subalgebra by $k\langle f_1, \dots, f_m\rangle$ or $k\langle \boldsymbol{f}\rangle$. As an example, we see that $t_1 \in \mathbb{R}\langle {t_1}^3\rangle$; more generally, if $(f_1, \dots, f_m) \in k[t_1, \dots, t_n]^m$ induces an injective map from k^n to k^m, we have $k\langle \boldsymbol{f}\rangle = k[t_1, \dots, t_n]$. In the present note, we will describe $k\langle \boldsymbol{f}\rangle$ in the case that k is algebraically closed and $\boldsymbol{f}$ induces a map from k^n to k^m that is surjective, or, in a sense specified later, at least close to being surjective.

The first set that $k\langle \boldsymbol{f}\rangle$ is compared with is the k-subalgebra of $k[t_1, \dots, t_n]$ generated by $\{f_1, \dots, f_m\}$, which we will denote by $k[f_1, \dots, f_m]$ or $k[\boldsymbol{f}]$; in this algebra, we find exactly those polynomials that can be written as $p(f_1, \dots, f_m)$ with $p \in k[x_1, \dots, x_m]$. Clearly, $k[\boldsymbol{f}] \subseteq k\langle \boldsymbol{f}\rangle$. The other inclusion need not hold in general: on any field k, let $f_1 = t_1$, $f_2 = t_1 t_2$. Then $\frac{f_2^2}{f_1} = t_1 t_2^2$ is (f_1, f_2)-determined, but $t_1 t_2^2 \notin k[f_1, f_2]$.

2010 *Mathematics Subject Classification.* 13B25 (12E05).

Key words and phrases. polynomial composition, polynomial maps.

Supported by the Austrian Science Fund (FWF):P24077.

The second set with which we will compare $k\langle \boldsymbol{f}\rangle$ is the set of all polynomials that can be written as rational functions in $f_1, \dots, f_m$. We denote the quotient field of $k[t_1, \dots, t_n]$ by $k(t_1, \dots, t_n)$. For $r_1, \dots, r_m \in k(t_1, \dots, t_n)$, the subfield of $k(t_1, \dots, t_n)$ that is generated by $k \cup \{r_1, \dots, r_m\}$ is denoted $k(r_1, \dots, r_m)$. We first observe that there are polynomials that can be written as rational functions in $\boldsymbol{f}$, but fail to be $\boldsymbol{f}$-determined. As an example, we see that $t_2 \in k(t_1, t_1t_2)$, but since $(0, 0\cdot 0) = (0, 0\cdot 1)$ and $0 \neq 1$, the polynomial t_2 is not (t_1, t_1t_2)-determined. As for the converse inclusion, we take a field k of positive characteristic χ. Then t_1 is $({t_1}^\chi)$-determined, but $t_1 \notin k({t_1}^\chi)$.

On the positive side, it is known that $k[f_1, \dots, f_m] = k\langle f_1, \dots, f_m\rangle$ holds in the following cases (cf. [1, Theorem 3.1]):

- k is algebraically closed, $m = n = 1$, and the derivative f' of f is not the zero polynomial, and, more generally,
- k is algebraically closed, $m = n$, and there are univariate polynomials $g_1, \dots, g_m \in k[t]$ with $g_1' \neq 0, \dots, g_m' \neq 0$, $f_1 = g_1(t_1), \dots, f_m = g_m(t_m)$.

Let us now briefly outline the results obtained in the present note: Let k be an algebraically closed field of characteristic 0, and let $f_1, \dots, f_m \in k[t_1, \dots, t_n]$ be algebraically independent over k. Then we have $k\langle \boldsymbol{f}\rangle \subseteq k(\boldsymbol{f})$ (Theorem 3.3). The equality $k[\boldsymbol{f}] = k\langle \boldsymbol{f}\rangle$ holds if and only if $\boldsymbol{f}$ induces a map from k^n to k^m that is *almost surjective* (see Definition 2.1). This equality is stated in Theorem 3.4. Similar results are given for the case of positive characteristic.

The last equality has a consequence on the functional decomposition of polynomials. If $\boldsymbol{f}$ induces a surjective mapping from k^n to k^m, (k algebraically closed of characteristic 0), and if $h : k^m \to k$ is an arbitrary function such that $h \circ \boldsymbol{f}$ is a polynomial function, then h is a polynomial function. In an algebraically closed field of positive characteristic χ, we will conclude that h is a composition of taking χ-th roots and a polynomial function (Corollary 4.2).

2. Preliminaries about polynomials

For the notions from algebraic geometry used in this note, we refer to [2]; deviating from their definitions, we call the set of solutions of a system of polynomial equations an *algebraic set* (instead of *affine variety*). For an algebraically closed field k and $A \subseteq k^m$, we let $I_m(A)$ (or simply $I(A)$) be the set of polynomials vanishing on every point in A, and for $P \subseteq k[t_1, \dots, t_m]$, we let $V_m(P)$ (or simply $V(P)$) be the set of common zeroes of P in k^m. The Zariski-closure $V(I(A))$ of a set $A \subseteq k^m$ will be abbreviated by $\overline{A}$. The *dimension* of an algebraic set A is the maximal $d \in \{0, \dots, m\}$ such that there are $i_1 < i_2 < \cdots < i_d \in \{1, \dots, m\}$ with $I(A) \cap k[x_{i_1}, \dots, x_{i_d}] = \{0\}$. We abbreviate the dimension of A by $\dim(A)$ and set $\dim(\varnothing) := -1$. For $f_1, \dots, f_m, g \in k[t_1, \dots, t_n]$, and

$D := \{(f_1(\boldsymbol{a}), \ldots, f_m(\boldsymbol{a}), g(\boldsymbol{a})) \mid \boldsymbol{a} \in k^n\}$, its Zariski-closure $\overline{D}$ is an irreducible algebraic set, and its dimension is the maximal number of algebraically independent elements in $\{f_1, \ldots, f_m, g\}$. The Closure Theorem [2, p. 258] tells that there exists an algebraic set $W \subseteq k^{m+1}$ with $\dim(W) < \dim(\overline{D})$ such that $\overline{D} = D \cup W$. If $\dim(\overline{D}) = m$, then there exists an irreducible polynomial $p \in k[x_1, \ldots, x_{m+1}]$ such that $\overline{D} = V(p)$. We will denote this p by $\mathrm{Irr}(\overline{D})$; $\mathrm{Irr}(\overline{D})$ is then defined up to a multiplication with a nonzero element from k. Above this, we recall that a set is *constructible* if and only if it can be generated from algebraic sets by a finite application of the set-theoretic operations of forming the union of two sets, the intersection of two sets, and the complement of a set, and that the range of a polynomial map from k^n to k^m and its complement are constructible. This is of course a consequence of the Theorem of Chevalley-Tarski [4, Exercise II.3.19], but since we are only concerned with the image of k^n, it also follows from [2, p.262, Corollary 2].

Definition 2.1. Let k be an algebraically closed field, $m, n \in \mathbb{N}$, and let $\boldsymbol{f} = (f_1, \ldots, f_m) \in (k[t_1, \ldots, t_n])^m$. By $\mathrm{range}(\boldsymbol{f})$, we denote the image of the mapping $\hat{\boldsymbol{f}} : k^n \to k^m$ that is induced by $\boldsymbol{f}$. We say that $\boldsymbol{f}$ is *almost surjective on* k if the dimension of the Zariski-closure of $k^m \setminus \mathrm{range}(\boldsymbol{f})$ is at most $m - 2$.

Proposition 2.2. *Let k be an algebraically closed field, and let $(f_1, \ldots, f_m) \in k[t_1, \ldots, t_n]^m$ be almost surjective on k. Then the sequence $(f_1, \ldots, f_m)$ is algebraically independent over k.*

Proof: Seeking a contradiction, we suppose that there is $u \in k[x_1, \ldots, x_m]$ with $u \neq 0$ and $u(f_1, \ldots, f_m) = 0$. Then $\mathrm{range}(\boldsymbol{f}) \subseteq V(u)$, hence $\dim(\overline{\mathrm{range}(\boldsymbol{f})}) \leq m - 1$. Since $\boldsymbol{f}$ is almost surjective, k^m is then the union of two algebraic sets of dimension $\leq m - 1$, a contradiction. □

We will use the following easy consequence of the description of constructible sets:

Proposition 2.3. *Let k be an algebraically closed field, and let B be a constructible subset of k^m with $\dim(\overline{B}) \geq m-1$. Then there exist algebraic sets W, X such that W is irreducible, $\dim(W) = m - 1$, $\dim(X) \leq m - 2$, $W \setminus X \subseteq B$.*

Since B is constructible, there are irreducible algebraic sets $V_1, \ldots, V_p$ and algebraic sets $W_1, \ldots, W_p$ with $W_i \subsetneq V_i$ and $B = \bigcup_{i=1}^{p}(V_i \setminus W_i)$ (cf. [2, p. 262]). We assume that the V_i's are ordered with nonincreasing dimension. If $\dim(V_1) = m$, then $k^m \setminus W_1 \subseteq B$. Let U be an irreducible algebraic set of dimension $m - 1$ with $U \not\subseteq W_1$. Then $U \cap (k^m \setminus W_1) = U \setminus (W_1 \cap U)$. Since $W_1 \cap U \neq U$, setting $W := U, X := W_1 \cap U$ yields the required sets.

If $\dim(V_1) = m - 1$, then $W := V_1$ and $X := W_1$ are the required sets.

The case $\dim(V_1) \leq m - 2$ cannot occur because then $\overline{B} \subseteq V_1 \cup \ldots \cup V_p$ has dimension at most $m - 2$. □

Let k be a field, and let $p, q, f \in k[t]$ such that $\deg(f) > 0$. It is known that $p(f)$ divides $q(f)$ if and only if p divides q [3, Lemma 2.1 and 2.2]. The following Lemma yields a multivariate version of this result.

Lemma 2.4. *Let k be an algebraically closed field, $m, n \in \mathbb{N}$, and let $\boldsymbol{f} = (f_1, \dots, f_m) \in (k[t_1, \dots, t_n])^m$. Then the following are equivalent:*

1. *$\boldsymbol{f}$ is almost surjective on k.*
2. *$k(f_1, \dots, f_m) \cap k[t_1, \dots, t_n] = k[f_1, \dots, f_m]$ and $(f_1, \dots, f_m)$ is algebraically independent over k.*
3. *For all $p, q \in k[x_1, \dots, x_m]$ with $p(f_1, \dots, f_m) \mid q(f_1, \dots, f_m)$, we have $p \mid q$.*

Proof: (1)⇒(2): (This proof uses some ideas from the proof of Theorem 4.2.1 in [5, p. 82].) Let $g \in k(f_1, \dots, f_m) \cap k[t_1, \dots, t_n]$. Then there are $r, s \in k[x_1, \dots, x_m]$ with $\gcd(r, s) = 1$ and $g = r(f_1, \dots, f_m)/s(f_1, \dots, f_m)$, and thus

$$g(t_1, \dots, t_n) \cdot s(f_1, \dots, f_m) = r(f_1, \dots, f_m). \tag{2.1}$$

Suppose $s \notin k$. Then $V(s)$ has dimension $m - 1$. We have $V(s) = (V(s) \cap \operatorname{range}(\boldsymbol{f})) \cup (V(s) \cap (k^m \setminus \operatorname{range}(\boldsymbol{f}))) \subseteq \overline{V(s) \cap \operatorname{range}(\boldsymbol{f})} \cup \overline{V(s) \cap (k^m \setminus \operatorname{range}(\boldsymbol{f}))}$. Since $\boldsymbol{f}$ is almost surjective, $\overline{V(s) \cap \operatorname{range}(\boldsymbol{f})}$ is then of dimension $m - 1$. Hence it contains an irreducible component of dimension $m - 1$, and thus there is an irreducible $p \in k[x_1, \dots, x_m]$ such that $V(p) \subseteq \overline{V(s) \cap \operatorname{range}(\boldsymbol{f})}$. Since then $V(p) \subseteq V(s)$, the Nullstellensatz yields $n_1 \in \mathbb{N}$ with $p \mid s^{n_1}$, and thus by the irreducibility of p, $p \mid s$. Now we show that for all $\boldsymbol{a} \in V(s) \cap \operatorname{range}(\boldsymbol{f})$, we have $r(\boldsymbol{a}) = 0$. To this end, let $\boldsymbol{b} \in k^n$ with $\boldsymbol{f}(\boldsymbol{b}) = \boldsymbol{a}$. Setting $\boldsymbol{t} := \boldsymbol{b}$ in (2.1), we obtain $r(\boldsymbol{a}) = 0$. Thus $V(s) \cap \operatorname{range}(\boldsymbol{f}) \subseteq V(r)$, and therefore $\overline{V(s) \cap \operatorname{range}(\boldsymbol{f})} \subseteq V(r)$, which implies $V(p) \subseteq V(r)$. By the Nullstellensatz, we have an $n_2 \in \mathbb{N}$ with $p \mid r^{n_2}$ and thus by the irreducibility of p, $p \mid r$. Now $p \mid r$ and $p \mid s$, contradicting $\gcd(r, s) = 1$. Hence $s \in k$, and thus $g \in k[f_1, \dots, f_m]$. The algebraic independence of $(f_1, \dots, f_m)$ follows from Proposition 2.2.

(2)⇒(3): Let $p, q \in k[x_1, \dots, x_m]$ such that $p(f_1, \dots, f_m) \mid q(f_1, \dots, f_m)$. If $p(f_1, \dots, f_m) = 0$, then $q(f_1, \dots, f_m) = 0$, and thus by the algebraic independence of $(f_1, \dots, f_m)$, we have $q = 0$ and thus $p \mid q$. Now assume $p(f_1, \dots, f_m) \neq 0$. We have $a(t_1, \dots, t_n) \in k[t_1, \dots, t_n]$ such that

$$q(f_1, \dots, f_m) = a(t_1, \dots, t_n) \cdot p(f_1, \dots, f_m) \tag{2.2}$$

and thus $a(t_1, \dots, t_n) \in k(f_1, \dots, f_m) \cap k[t_1, \dots, t_n]$. Thus there exists $b \in k[x_1, \dots, x_m]$ such that $a(t_1, \dots, t_n) = b(f_1, \dots, f_m)$. Now (2.2) yields

$$q(f_1, \dots, f_m) = b(f_1, \dots, f_m) \cdot p(f_1, \dots, f_m).$$

Using the algebraic independence of $(f_1, \dots, f_m)$, we obtain $q(x_1, \dots, x_m) = b(x_1, \dots, x_m) \cdot p(x_1, \dots, x_m)$, and thus $p \mid q$.

(3)⇒(1) Seeking a contradiction, we suppose that $\boldsymbol{f}$ is not almost surjective. Let $B := k^m \setminus \mathrm{range}(\boldsymbol{f})$. Then $\dim(\overline{B}) \geq m-1$. Since B is constructible, Proposition 2.3 yields W, X with W irreducible, $\dim(W) = m-1$, $\dim(X) \leq m-2$, $W \setminus X \subseteq B$. Since W is irreducible of dimension $m-1$, there is $p \in k[x_1, \ldots, x_m]$ such that $W = V(p)$. Since $\dim(W) > \dim(X)$, we have $W \not\subseteq X$, thus $I(X) \not\subseteq I(W)$, and therefore there is $q \in I(X)$ with $q \notin I(W)$. We have $W \subseteq B \cup X$, and thus $W \cap \mathrm{range}(\boldsymbol{f}) \subseteq X$. This implies that for all $\boldsymbol{a} \in k^n$ with $p(\boldsymbol{f}(\boldsymbol{a})) = 0$, we have $q(\boldsymbol{f}(\boldsymbol{a})) = 0$: in fact, if $p(\boldsymbol{f}(\boldsymbol{a})) = 0$, then $\boldsymbol{f}(\mathbf{a}) \in V(p) \cap \mathrm{range}(\boldsymbol{f}) = W \cap \mathrm{range}(\boldsymbol{f}) \subseteq X$. Hence $q(\boldsymbol{f}(\boldsymbol{a})) = 0$. By the Nullstellensatz, we obtain a $\nu \in \mathbb{N}$ such that $p(f_1, \ldots, f_m) \mid q(f_1, \ldots, f_m)^\nu$. Therefore, using (3), we have $p \mid q^\nu$. This implies $V(p) \subseteq V(q)$. Thus we have $W \subseteq V(q)$ and therefore $q \in I(W)$, contradicting the choice of q. Hence $\boldsymbol{f}$ is almost surjective, proving (1). □

3. $\boldsymbol{f}$-determined polynomials

We will first show that often all $\boldsymbol{f}$-determined polynomials are rational functions of $\boldsymbol{f}$. Special care, however, is needed in the case of positive characteristic. In an algebraically closed field of characteristic $\chi > 0$, the unary polynomial t_1 is (t_1^χ)-determined, but t_1 is neither a polynomial nor a rational function of t_1^χ.

Definition 3.1. Let k be a field of characteristic $\chi > 0$, let $n \in \mathbb{N}$, and let P be a subset of $k[t_1, \ldots, t_n]$. We define the set $\mathrm{rad}_\chi(P)$ by

$$\mathrm{rad}_\chi(P) := \{f \in k[t_1, \ldots, t_n] \mid \text{ there is } \nu \in \mathbb{N}_0 \text{ such that } f^{\chi^\nu} \in P\}.$$

Lemma 3.2. *Let k be an algebraically closed field, let $m, n \in \mathbb{N}$, let $f_1, \ldots, f_m$ be algebraically independent polynomials in $k[t_1, \ldots, t_n]$, let $g \in k\langle f_1, \ldots, f_m\rangle$, and let $D := \{(f_1(\boldsymbol{a}), \ldots, f_m(\boldsymbol{a}), g(\boldsymbol{a})) \mid \boldsymbol{a} \in k^n\}$. Then $\dim(\overline{D}) = m$.*

Proof: By the closure theorem [2, p. 258], there is an algebraic set W such that $\overline{D} = D \cup W$ and $\dim(W) < \dim(\overline{D})$. Let $\pi : k^{m+1} \to k^m, (y_1, \ldots, y_{m+1}) \mapsto (y_1, \ldots, y_m)$ be the projection of k^{m+1} onto the first m coordinates, and let $\overline{\pi(W)}$ be the Zariski-closure of $\pi(W)$ in k^m. We will now examine the projection of D. Since $(f_1, \ldots, f_m)$ is algebraically independent, $\pi(D)$ is Zariski-dense in k^m, and hence $\dim(\overline{\pi(D)}) = m$. Since $\dim(V) \geq \dim(\overline{\pi(V)})$ holds for every algebraic set V, we then obtain $\dim(\overline{D}) \geq \dim(\overline{\pi(\overline{D})}) \geq \dim(\overline{\pi(D)}) = m$. Seeking a contradiction, we suppose that $\dim(\overline{D}) = m+1$.

In the case $\dim(\overline{\pi(W)}) = m$, we use [2, p. 193, Theorem 3], which tells $\overline{\pi(W)} = V_m(I(W) \cap k[x_1, \ldots, x_m])$, and we obtain that $k^m = V_m(I(W) \cap k[x_1, \ldots, x_m])$, and therefore $I(W) \cap k[x_1, \ldots, x_m] = \{0\}$. Hence, $x_1 + I(W), \ldots, x_m + I(W)$ are algebraically independent in $k[x_1, \ldots, x_{m+1}]/I(W)$. Since $\dim(W) \leq m$, we observe that the sequence $(x_1 + I(W), \ldots, x_{m+1} + I(W))$ is algebraically dependent over k, and therefore, there is a polynomial $q(x_1, \ldots, x_{m+1}) \in I(W)$ with

$\deg_{x_{m+1}}(q) > 0$. Let r be the leading coefficient of q with respect to x_{m+1}, and let $(y_1, \dots, y_m) \in k^m$ be such that $r(y_1, \dots, y_m) \neq 0$. Then there are only finitely many $z \in k$ with $(y_1, \dots, y_m, z) \in W$. Since $\overline{D} = k^{m+1}$, there are then infinitely many $z \in k$ with $(y_1, \dots, y_m, z) \in D$, a contradiction to the fact that g is $\boldsymbol{f}$-determined.

In the case $\dim(\overline{\pi(W)}) \leq m-1$, we take $(y_1, \dots, y_m) \in k^m \setminus \pi(W)$. For all $z \in k$, we have $(y_1, \dots, y_m, z) \in \overline{D}$ and $(y_1, \dots, y_m, z) \notin W$, and therefore all $(y_1, \dots, y_m, z)$ are elements of D, a contradiction to the fact that g is $\boldsymbol{f}$-determined.

Hence, we have $\dim(\overline{D}) = m$. □

Theorem 3.3. *Let k be an algebraically closed field, let χ be its characteristic, let $m, n \in \mathbb{N}$, and let $(f_1, \dots, f_m)$ be a sequence of polynomials in $k[t_1, \dots, t_n]$ that is algebraically independent over k. Then we have:*

(1) *If $\chi = 0$, then $k\langle f_1, \dots, f_m\rangle \subseteq k(f_1, \dots, f_m) \cap k[t_1, \dots, t_n]$.*
(2) *If $\chi > 0$, then $k\langle f_1, \dots, f_m\rangle \subseteq \mathrm{rad}_\chi(k(f_1, \dots, f_m) \cap k[t_1, \dots, t_n])$.*

Proof: Let $g \in k\langle f_1, \dots, f_m\rangle$. We define

$$D := \{(f_1(\boldsymbol{a}), \dots, f_m(\boldsymbol{a}), g(\boldsymbol{a})) \mid \boldsymbol{a} \in k^n\},$$

we let $\overline{D}$ be its Zariski-closure in k^{m+1}, and we let W be an algebraic set with $\dim(W) < \dim(\overline{D})$ and $\overline{D} = D \cup W$. By Lemma 3.2, we have $\dim(\overline{D}) = m$. Now, we distinguish cases according to the characteristic of k. Let us first suppose $\chi = 0$. Let $q := \mathrm{Irr}(\overline{D})$ be an irreducible polynomial with $\overline{D} = V(q)$, and let $d := \deg_{x_{m+1}}(q)$. Since $f_1, \dots, f_m$ are algebraically independent over k, we have $d \geq 1$. We will now prove $d = 1$. Suppose $d > 1$. We write $q = \sum_{i=0}^{d} q_i(x_1, \dots, x_m)\, x_{m+1}^i$. We recall that for a field K, and $f, g \in K[t]$ of positive degree, the resultant $\mathrm{res}_t(f, g)$ is 0 if and only if $\deg(\gcd_{K[t]}(f, g)) \geq 1$ [2, p. 156, Proposition 8]. Let $r := \mathrm{res}_{x_{m+1}}(q, \frac{\partial}{\partial x_{m+1}} q)$ be the resultant of q and its derivative when seen as elements of the ring $k(x_1, \dots, x_m)[x_{m+1}]$. If $r = 0$, then q and $\frac{\partial}{\partial x_{m+1}} q$ have a common divisor in $k(x_1, \dots, x_m)[x_{m+1}]$ with $1 \leq \deg_{x_{m+1}}(q) \leq d-1$ in $k(x_1, \dots, x_m)[x_{m+1}]$. Using a standard argument involving Gauß's Lemma, we find a divisor a of q in $k[x_1, \dots, x_{m+1}]$ such that $1 \leq \deg_{x_{m+1}}(a) \leq d-1$. This contradicts the irreducibility of q. Hence $r \neq 0$. Since $\dim(\overline{\pi(W)}) \leq m-1$, $r \neq 0$, and $q_d \neq 0$, we have $V(r) \cup V(q_d) \cup \pi(W) \neq k^m$. Thus we can choose $\boldsymbol{a} \in k^m$ such that $r(\boldsymbol{a}) \neq 0$, $q_d(\boldsymbol{a}) \neq 0$, and $\boldsymbol{a} \notin \pi(W)$. Let $\tilde{q}(t) := q(\boldsymbol{a}, t)$. Since $\mathrm{res}_t(\tilde{q}(t), \tilde{q}'(t)) = r(\boldsymbol{a}) \neq 0$, $\tilde{q}$ has d different roots in k, and thus $q(\boldsymbol{a}, x) = 0$ has d distinct solutions for x, say $b_1, \dots, b_d$. We will now show $\{(\boldsymbol{a}, b_i) \mid i \in \{1, \dots, d\}\} \subseteq D$. Let $i \in \{1, \dots, d\}$, and suppose that $(\boldsymbol{a}, b_i) \notin D$. Then $(\boldsymbol{a}, b_i) \in W$, and thus $\boldsymbol{a} \in \pi(W)$, a contradiction. Thus all the elements $(\boldsymbol{a}, b_1), \dots, (\boldsymbol{a}, b_d)$ lie in D. Since $d > 1$, this implies that g is not

$(f_1, \ldots, f_m)$-determined. Therefore we have $d = 1$. Since $(f_1, \ldots, f_m)$ is algebraically independent, the polynomial q witnesses that g is algebraic of degree 1 over $k(f_1, \ldots, f_m)$, and thus lies in $k(f_1, \ldots, f_m)$. This concludes the case $\chi = 0$.

Now we assume $\chi > 0$. It follows from Lemma 3.2 that for every $h \in k\langle t_1, \ldots, t_n\rangle$, the Zariski-closure of $D(h) := \{(f_1(\boldsymbol{a}), \ldots, f_m(\boldsymbol{a}), h(\boldsymbol{a})) \mid \boldsymbol{a} \in k^n\}$ is an irreducible variety of dimension m in k^{m+1}. This implies that there is an irreducible polynomial $\mathrm{Irr}(\overline{D(h)}) \in k[x_1, \ldots, x_m]$ such that $\overline{D(h)} = V(\mathrm{Irr}(\overline{D(h)}))$. Furthermore, by the closure theorem [2], there is an algebraic set $W(h) \subseteq k^m$ such that $\dim(W(h)) \leq m - 1$ and $D(h) \cup W(h) = \overline{D(h)}$. We will now prove the following statement by induction on $\deg_{x_{m+1}}(\mathrm{Irr}(\overline{D(h)}))$.

> Every $\boldsymbol{f}$-determined polynomial $h \in k[t_1, \ldots, t_n]$ is an element of $\mathrm{rad}_\chi(k(f_1, \ldots, f_m) \cap k[t_1, \ldots, t_n])$.

Let

$$d := \deg_{x_{m+1}}(\mathrm{Irr}(\overline{D(h)})).$$

If $d = 0$, then $f_1, \ldots, f_m$ are algebraically dependent, a contradiction. If $d = 1$, then since $f_1, \ldots, f_m$ are algebraically independent, h is algebraic of degree 1 over $k(f_1, \ldots, f_m)$, and thus lies in $k(f_1, \ldots, f_m) \cap k[t_1, \ldots, t_n]$. Let us now consider the case $d > 1$. We set

$$e := \deg_{x_{m+1}}(\tfrac{\partial}{\partial x_{m+1}}\mathrm{Irr}(\overline{D(h)})).$$

If $\frac{\partial}{\partial x_{m+1}}\mathrm{Irr}(\overline{D(h)}) = 0$, then there is a polynomial $p \in k[x_1, \ldots, x_{m+1}]$ such that $\mathrm{Irr}(\overline{D(h)}) = p(x_1, \ldots, x_m, x_{m+1}^\chi)$. We know that h^χ is $\boldsymbol{f}$-determined, hence by Lemma 3.2, $\overline{D(h^\chi)}$ is of dimension m. Since

$$p\,(f_1, \ldots, f_m, h^\chi) = \mathrm{Irr}(\overline{D(h)})\,(f_1, \ldots, f_m, h) = 0,$$

we have $p \in I(D(h^\chi))$. Thus $\overline{D(h^\chi)} \subseteq V(p)$. Therefore, the irreducible polynomial $\mathrm{Irr}(\overline{D(h^\chi)})$ divides p, and thus

$$\deg_{x_{m+1}}(\mathrm{Irr}(\overline{D(h^\chi)})) \leq \deg_{x_{m+1}}(p) < \deg_{x_{m+1}}(\mathrm{Irr}(\overline{D(h)})).$$

By the induction hypothesis, we obtain that h^χ is an element of $\mathrm{rad}_\chi(k(f_1, \ldots, f_m) \cap k[t_1, \ldots, t_n])$. Therefore, $h \in \mathrm{rad}_\chi(k(f_1, \ldots, f_m) \cap k[t_1, \ldots, t_n])$. This concludes the case that $\frac{\partial}{\partial x_{m+1}}(\mathrm{Irr}(\overline{D(h)})) = 0$.

If $e = 0$, we choose $\boldsymbol{a} = (a_1, \ldots, a_m) \in k^m$ such that

$$\frac{\partial}{\partial x_{m+1}}\mathrm{Irr}(\overline{D(h)})\ (a_1, \ldots, a_m, 0) \neq 0,$$

such that the leading coefficient of $\mathrm{Irr}(\overline{D(h)})$ with respect to x_{m+1} does not vanish at $\boldsymbol{a}$, and such that $\boldsymbol{a} \notin \pi(W(h))$. Then $\mathrm{Irr}(\overline{D(h)})(\boldsymbol{a}, x) = 0$ has d different solutions for x, say $b_1, \ldots, b_d$. Since $\{(\boldsymbol{a}, b_i) \mid i \in \{1, \ldots, d\}\} \cap W(h) = \varnothing$ because

$\boldsymbol{a} \notin \pi(W(h))$, we have $\{(\boldsymbol{a}, b_i) \mid i \in \{1, \dots, d\}\} \subseteq D(h)$. Since h is $\boldsymbol{f}$-determined, $d = 1$, contradicting the case assumption.

If $e > 0$, then we compute the resultant $r := \operatorname{res}^{(d,e)}_{x_{m+1}}(\operatorname{Irr}(\overline{D(h)}), \frac{\partial}{\partial x_{m+1}}\operatorname{Irr}(\overline{D(h)}))$, seen as polynomials of degrees d and e over the field $k(x_1, \dots, x_m)$ in the variable x_{m+1}. As in the case $\chi = 0$, the irreducibility of $\operatorname{Irr}(\overline{D(h)})$ yields $r \neq 0$. Now we let $\boldsymbol{a} \in k^m$ be such that $r(\boldsymbol{a}) \neq 0$, the leading coefficient $(\operatorname{Irr}(\overline{D(h)}))_d$ of $\operatorname{Irr}(\overline{D(h)})$ with respect to x_{m+1} does not vanish at $\boldsymbol{a}$, and $\boldsymbol{a} \notin \pi(W(h))$. Setting $\tilde{q}(t) := \operatorname{Irr}(\overline{D(h)})\,(\boldsymbol{a}, t)$, we see that $\operatorname{res}^{(d,e)}_t(\tilde{q}(t), \tilde{q}'(t)) \neq 0$. Thus $\tilde{q}$ has d distinct zeroes $b_1, \dots, b_d$, and then $\{(\boldsymbol{a}, b_i) \mid i \in \{1, \dots, d\}\} \subseteq D(h)$. Since $d > 1$, this contradicts the fact that h is $\boldsymbol{f}$-determined. □

Theorem 3.4. *Let k be an algebraically closed field of characteristic 0, let $m, n \in \mathbb{N}$, and let $\boldsymbol{f} = (f_1, \dots, f_m)$ be a sequence of algebraically independent polynomials in $k[t_1, \dots, t_n]$. Then the following are equivalent:*

(1) $k\langle f_1, \dots, f_m\rangle = k[f_1, \dots, f_m]$.
(2) $\boldsymbol{f}$ *is almost surjective.*

Proof: (1)⇒(2): Suppose that $\boldsymbol{f}$ is not almost surjective. Then by Lemma 2.4, there are $p, q \in k[x_1, \dots, x_m]$ such that $p(f_1, \dots, f_m) \mid q(f_1, \dots, f_m)$ and $p \nmid q$. Let $d := \gcd(p, q)$, $p_1 := p/d$, $q_1 := q/d$. Let $a(t_1, \dots, t_n) \in k[t_1, \dots, t_n]$ be such that

$$p_1(f_1, \dots, f_m) \cdot a(t_1, \dots, t_n) = q_1(f_1, \dots, f_m). \tag{3.1}$$

We claim that $b(t_1, \dots, t_n) := q_1(f_1, \dots, f_m) \cdot a(t_1, \dots, t_m)$ is $\boldsymbol{f}$-determined and is not an element of $k[f_1, \dots, f_m]$. In order to show that b is $\boldsymbol{f}$-determined, we let $\boldsymbol{c}, \boldsymbol{d} \in k^n$ such that $\boldsymbol{f}(\boldsymbol{c}) = \boldsymbol{f}(\boldsymbol{d})$. If $p_1(\boldsymbol{f}(\boldsymbol{c})) \neq 0$, we have $b(\boldsymbol{c}) = q_1(\boldsymbol{f}(\boldsymbol{c})) \cdot a(\boldsymbol{c}) = q_1(\boldsymbol{f}(\boldsymbol{c})) \cdot \frac{q_1(\boldsymbol{f}(\boldsymbol{c}))}{p_1(\boldsymbol{f}(\boldsymbol{c}))} = q_1(\boldsymbol{f}(\boldsymbol{d})) \cdot \frac{q_1(\boldsymbol{f}(\boldsymbol{d}))}{p_1(\boldsymbol{f}(\boldsymbol{d}))} = q_1(\boldsymbol{f}(\boldsymbol{d})) \cdot a(\boldsymbol{d}) = b(\boldsymbol{d})$. If $p_1(\boldsymbol{f}(\boldsymbol{c})) = 0$, we have $b(\boldsymbol{c}) = q_1(\boldsymbol{f}(\boldsymbol{c})) \cdot a(\boldsymbol{c})$. By (3.1), we have $q_1(\boldsymbol{f}(\boldsymbol{c})) = 0$, and thus $b(\boldsymbol{c}) = 0$. Similarly $b(\boldsymbol{d}) = 0$. This concludes the proof that b is $\boldsymbol{f}$-determined.

Let us now show that $b \notin k[f_1, \dots, f_m]$. We have

$$b(t_1, \dots, t_n) = \frac{q_1(f_1, \dots, f_m)^2}{p_1(f_1, \dots, f_m)}.$$

If $b \in k[f_1, \dots, f_m]$, there is $r \in k[x_1, \dots, x_m]$ with $r(f_1, \dots, f_m) = b(t_1, \dots, t_n)$. Then $r(f_1, \dots, f_m) \cdot p_1(f_1, \dots, f_m) = q_1(f_1, \dots, f_m)^2$. From the algebraic independence of $(f_1, \dots, f_m)$, we obtain $r(x_1, \dots, x_m) \cdot p_1(x_1, \dots, x_m) = q_1(x_1, \dots, x_m)^2$, hence $p_1(x_1, \dots, x_m) \mid q_1(x_1, \dots, x_m)^2$. Since p_1, q_1 are relatively prime, we then have $p_1(x_1, \dots, x_m) \mid q_1(x_1, \dots, x_m)$, contradicting the choice of p and q. Hence $\boldsymbol{f}$ is almost surjective.

(2)⇒(1): From Theorem 3.3, we obtain $k\langle \boldsymbol{f}\rangle \subseteq k(\boldsymbol{f}) \cap k[t_1,\ldots,t_n]$. Since $\boldsymbol{f}$ is almost surjective, Lemma 2.4 yields $k(\boldsymbol{f}) \cap k[t_1,\ldots,t_n] = k[\boldsymbol{f}]$, and thus $k\langle \boldsymbol{f}\rangle \subseteq k[\boldsymbol{f}]$. The other inclusion is obvious. □

Theorem 3.5. *Let k be an algebraically closed field of characteristic $\chi > 0$, let $m, n \in \mathbb{N}$, and let $\boldsymbol{f} = (f_1,\ldots,f_m)$ be a sequence of algebraically independent polynomials in $k[t_1,\ldots,t_n]$. Then the following are equivalent:*

(1) $k\langle f_1,\ldots,f_m\rangle = \mathrm{rad}_\chi(k[f_1,\ldots,f_m])$.
(2) $\boldsymbol{f}$ *is almost surjective.*

Proof: (1)⇒(2): As in the proof of Theorem 3.4, we produce an $\boldsymbol{f}$-determined polynomial b and relatively prime $p_1, q_1 \in k[x_1,\ldots,x_m]$ with $p_1 \nmid q_1$ and

$$b(t_1,\ldots,t_n) = \frac{q_1(f_1,\ldots,f_m)^2}{p_1(f_1,\ldots,f_m)}.$$

Now suppose that there is $\nu \in \mathbb{N}_0$ with $b^{\chi^\nu} \in k[f_1,\ldots,f_m]$. Then $p_1(f_1,\ldots,f_m)^{\chi^\nu}$ divides
$q_1(f_1,\ldots,f_m)^{2\chi^\nu}$ in $k[f_1,\ldots,f_m]$, and thus $p_1(x_1,\ldots,x_m)$ divides $q_1(x_1,\ldots,x_m)^{2\chi^\nu}$ in $k[x_1,\ldots,x_m]$. Since p_1 and q_1 are relatively prime, we obtain $p_1 \mid q_1$, contradicting the choice of p_1 and q_1.

(1)⇒(2): From Theorem 3.3, we obtain $k\langle \boldsymbol{f}\rangle \subseteq \mathrm{rad}_\chi(k(\boldsymbol{f}) \cap k[t_1,\ldots,t_n])$. Since $\boldsymbol{f}$ is almost surjective, Lemma 2.4 yields $k(\boldsymbol{f}) \cap k[t_1,\ldots,t_n] = k[\boldsymbol{f}]$, and thus $k\langle \boldsymbol{f}\rangle \subseteq \mathrm{rad}_\chi(k[\boldsymbol{f}])$. The other inclusion follows from the fact that the map $\varphi : k \to k$, $\varphi(y) := y^\chi$ is injective. □

4. Function compositions that are polynomials

For a field k, let $\boldsymbol{f} = (f_1,\ldots,f_m) \in (k[t_1,\ldots,t_n])^m$, and let $h : k^m \to k$ be an arbitrary function. Then we write $h \circ \boldsymbol{f}$ for the function defined by $(h \circ \boldsymbol{f})(\boldsymbol{a}) = h(f_1(\boldsymbol{a}),\ldots,f_m(\boldsymbol{a}))$ for all $\boldsymbol{a} \in k^n$. For an algebraically closed field K of characteristic $\chi > 0$, $y \in K$ and $\nu \in \mathbb{N}_0$, we let $s^{(\chi^\nu)}(y)$ be the element in K with $(s^{(\chi^\nu)}(y))^{\chi^\nu} = y$; so $s^{(\chi^\nu)}$ takes the χ^ν-th root.

Theorem 4.1. *Let k be a field, let K be its algebraic closure, let $m, n \in \mathbb{N}$, let $g, f_1,\ldots,f_m \in k[t_1,\ldots,t_n]$, and let $h : K^m \to K$ be an arbitrary function. Let $R := \boldsymbol{f}(K^n)$ be the range of the function from K^n to K^m that $\boldsymbol{f} = (f_1,\ldots,f_m)$ induces on K. We assume that $\dim(\overline{K^m \setminus R}) \leq m-2$, and that $h \circ \boldsymbol{f} = g$ on K, which means that*

$$h(\boldsymbol{f}(\boldsymbol{a})) = g(\boldsymbol{a}) \text{ for all } \boldsymbol{a} \in K^n.$$

Then we have:

(1) *If k is of characteristic 0, then there is $p \in k[x_1,\ldots,x_m]$ such that $h(\boldsymbol{b}) = p(\boldsymbol{b})$ for all $\boldsymbol{b} \in R$.*

(2) *If k is of characteristic $\chi > 0$, then there is $p \in k[x_1, \dots, x_m]$ and $\nu \in \mathbb{N}_0$ such that $h(\boldsymbol{b}) = s^{(\chi^\nu)}(p(\boldsymbol{b}))$ for all $\boldsymbol{b} \in R$.*

Proof: Let us first assume that k is of characteristic 0. We observe that as a polynomial in $K[t_1, \dots, t_n]$, g is $\boldsymbol{f}$-determined. Hence by Theorem 3.4, there is $q \in K[x_1, \dots, x_m]$ such that $q(f_1, \dots, f_m) = g$. Writing

$$q = \sum_{(i_1,\dots,i_m)\in I} \alpha_{i_1,\dots,i_m} x_1^{i_1} \cdots x_m^{i_m},$$

we obtain $g = \sum_{(i_1,\dots,i_m)\in I} \alpha_{i_1,\dots,i_m} f_1^{i_1} \cdots f_m^{i_m}$. Expanding the right hand side and comparing coefficients, we see that $(\alpha_{i_1,\dots,i_m})_{(i_1,\dots,i_m)\in I}$ is a solution of a linear system with coefficients in k. Since this system has a solution over K, it also has a solution over k. The solution over k provides the coefficients of a polynomial $p \in k[x_1, \dots, x_m]$ such that $p(f_1, \dots, f_m) = g$. From this, we obtain that $p(f_1(\boldsymbol{a}), \dots, f_m(\boldsymbol{a})) = g(\boldsymbol{a})$ for all $\boldsymbol{a} \in K^n$, and thus $p(\boldsymbol{b}) = h(\boldsymbol{b})$ for all $\boldsymbol{b} \in R$. This completes the proof of item (1).

In the case that k is of characteristic $\chi > 0$, Theorem 3.5 yields a polynomial $q \in K[x_1, \dots, x_m]$ and $\nu \in \mathbb{N}_0$ such that $q(f_1, \dots, f_m) = g^{\chi^\nu}$. As in the previous case, we obtain $p \in k[x_1, \dots, x_m]$ such that $p(f_1, \dots, f_m) = g^{\chi^\nu}$. Let $\boldsymbol{b} \in R$, and let $\boldsymbol{a}$ be such that $\boldsymbol{f}(\boldsymbol{a}) = \boldsymbol{b}$. Then $s^{(\chi^\nu)}(p(\boldsymbol{b})) = s^{(\chi^\nu)}(p(\boldsymbol{f}(\boldsymbol{a}))) = g(\boldsymbol{a}) = h(\boldsymbol{f}(\boldsymbol{a})) = h(\boldsymbol{b})$, which completes the proof of (2). □

We will now state the special case that k is algebraically closed and $\boldsymbol{f}$ is surjective in the following corollary. By a *polynomial function*, we will simply mean a function induced by a polynomial with all its coefficients in k.

Corollary 4.2. *Let k be an algebraically closed field, let $\boldsymbol{f} = (f_1, \dots, f_m) \in (k[t_1, \dots, t_n])^m$, and let $h : k^m \to k$ be an arbitrary function. We assume that $\boldsymbol{f}$ induces a surjective mapping from k^n to k^m and that $h \circ \boldsymbol{f}$ is a polynomial function. Then we have:*

(1) *If k is of characteristic 0, then h is a polynomial function.*
(2) *If k is of characteristic $\chi > 0$, then there is $\nu \in N_0$ such that $h^{\chi^\nu} : (y_1, \dots, y_m) \mapsto h(y_1, \dots, y_m)^{\chi^\nu}$ is a polynomial function.*

References

1. E. Aichinger and S. Steinerberger, *A proof of a theorem by Fried and MacRae and applications to the composition of polynomial functions*, Arch. Math. (Basel) **97** (2011), no. 2, 115–124.
2. D. Cox, J. Little, and D. O'Shea, *Ideals, varieties, and algorithms*, third ed., Undergraduate Texts in Mathematics, Springer, New York, 2007.
3. H. T. Engstrom, *Polynomial substitutions*, Amer. J. Math. **63** (1941), 249–255.
4. R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, no. 52, Springer-Verlag, New York, 1977.

5. A. van den Essen, *Polynomial automorphisms and the Jacobian conjecture*, Progress in Mathematics, vol. 190, Birkhäuser Verlag, Basel, 2000.

Erhard Aichinger, Institut für Algebra, Johannes Kepler Universität Linz, 4040 Linz, Austria

E-mail address: erhard@algebra.uni-linz.ac.at

URL: http://www.algebra.uni-linz.ac.at